

**Neomycin Sulfate and Oxytetracycline Growing Cattle (Cattle Over 400 lbs.) C
Type C Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATIONS FOR USE

For the reduction of the incidence of liver abscesses in growing cattle over 400 lbs.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate.....
grams/ton*
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to _____grams oxytetracycline hydrochloride/ton.* See FEEDING DIRECTIONS.

GUARANTEED ANALYSIS

Crude Protein, (Min).....XX.XX%
NPN, (Max)¹..... X.XX%
Crude Fat, (Min)..... X.XX%
Crude Fiber, (Max)..... X.XX%
Calcium, (Min)..... X.XX%
Calcium, (Max)..... X.XX%
Phosphorus, (Min)..... X.XX%
Salt, (Min)¹..... X.XX%
Salt, (Max)¹..... X.XX%
Sodium, (Min)²..... X.XX%
Sodium, (Max)²..... X.XX%
Potassium, (Min)¹..... X.XX%
Vitamin A, (Min)³..... ____IU/LB

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³Other than precursor of Vitamin A, if added

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed at 75 mg/head/day continuously as sole ration. For example, the table below with the stated drug concentration, body weight and feed consumption will provide cattle with 75 mg/head/day dosage.

Concentration of Oxytetracycline and Neomycin Sulfate (g/ton) in Type C Feed	Body Weight (lb)	Type C Feed Consumption (lb)
15	500	10
12	625	12.5
10	750	15

RESIDUE WARNING: Zero-day withdrawal period. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residue.

*The final printed medicated feed label must state a single drug concentration.

Net Weight _____lb. (_____kg.)

Manufactured by
Blue Bird Feed Company

Blue Bird, MD 00000