

**Neomycin Sulfate and Oxytetracycline Chicken Ration B
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD)
drug to use by or on the order of a licensed veterinarian.**

INDICATIONS FOR USE

For control of infectious synovitis caused by Mycoplasma synoviae; control of fowl cholera caused by Pasteurella multocida susceptible to oxytetracycline.

ACTIVE DRUG INGREDIENTS

Neomycin Sulfate.....100-200 grams/ton.*
Oxytetracycline (from oxytetracycline dihydrate base) equivalent to 100-200 grams
oxytetracycline hydrochloride/ton.*

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX%
Lysine, (Min).....	X.XX%
Methionine, (Min).....	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min).....	X.XX%
Calcium, (Max).....	X.XX%
Phosphorus, (Min).....	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by
the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO.

FEEDING DIRECTIONS

Feed continuously for 7-14 days as the sole ration.

RESIDUE WARNING: Do not feed to chickens producing eggs for human consumption.
Zero-day withdrawal period. In low calcium feeds, withdraw 3 days prior to slaughter.

*The final printed medicated feed label must state a single drug concentration.

Net Weight _____ lb. (_____ kg.)

Manufactured by
Blue Bird Feed Company
Blue bird, MD 00000