

**Chlortetracycline Turkey Ration II
Type C Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For the control of hexamitiasis caused by *Hexamita meleagridis* susceptible to chlortetracycline.

ACTIVE DRUG INGREDIENT

Chlortetracycline.....400 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than.....%
Crude Fat, not less than.....%
Crude Fiber, not more than.....%
Lysine (min).....%
Methionine (min).....%
Calcium (min).....%
Calcium (max).....%
Phosphorus(min).....%
Salt¹ (min).....%
Salt¹ (max).....%
Sodium² (min).....%
Sodium² (max).....%

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

INGREDIENTS

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

Feed continuously for 7 to 14 days as the sole ration.

WARNING: Do not feed to turkeys producing eggs for human consumption.
Zero-day withdrawal period.

Net Weight _____ lb. (_____ kg.)

Manufactured by
Blue Bird Feed Company
Blue bird, MD 00000