

**DCX + Chlortetracycline-Pennchlor Type B Cattle Feed**  
**Medicated Type B Medicated Feed**  
(decoquinatate and chlortetracycline Type B medicated feed)

**Caution:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

Calves, beef and non-lactating dairy cattle. For the prevention of coccidiosis caused by *Eimeria bovis* and *E. zuernii*; for treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* organisms susceptible to chlortetracycline.

**ACTIVE DRUG INGREDIENTS\***

Decoquinatate<sup>a</sup>.....535.8 to 1818 g/ton  
Chlortetracycline<sup>b</sup> (as hydrochloride)..... 20,000 to 80,000 g/ton

**GUARANTEED ANALYSIS**

Crude protein, not less than ..... %  
Crude fat, not less than ..... %  
Crude fiber, not more than ..... %  
Calcium, not less than ..... %  
Calcium, not more than ..... %  
Phosphorus, not less than ..... %  
Potassium, not less than ..... %  
Salt<sup>1</sup>, not less than ..... %  
Salt<sup>1</sup>, not more than ..... %  
Sodium<sup>2</sup>, not less than ..... %  
Sodium<sup>2</sup>, not more than ..... %

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials. This includes trace minerals and vitamins.

**MIXING DIRECTIONS**

In order to treat cattle for bacterial enteritis and/or pneumonia while preventing coccidiosis, mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of complete cattle feed containing 12.9 to 90.8 grams decoquinatate and 500 to 4000 grams chlortetracycline. The resulting Type C medicated feed should be fed at a rate to provide 1g chlortetracycline per 100 lb body weight/day and 22.7 mg decoquinatate per 100 lb of body weight/day daily for not more than 5 days. When this Type C medicated feed is fully consumed, resume feeding 22.7 mg decoquinatate per 100 lb of body weight/day for a total of 28 days to prevent coccidiosis. The following table provides examples of mixing rates.

Type B, grams/ton		Lbs Type B per Ton of Feed	Lbs Non-Medicated Feed Per Ton	Type C, grams/ton	
Decoquinatate	CTC			Decoquinatate	CTC
568	25,000	80	1920	22.7	1000
1136	50,000	40	1960	22.7	1000
1136	50,000	400	1600	227	10,000
1818	80,000	250	1750	227	10,000

**Caution:** Bentonite should not be used in decoquinatate feeds.



**Warning:** Withdraw 24 hours prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Don't use in calves to be processed for veal. Do not feed to animals producing milk for food.



Lot Number (if applicable): \_\_\_\_\_

**MANUFACTURED BY**

Blue Bird Feed Company  
Robin, Indiana 11111

**NET WEIGHT ON BAG OR BULK**

\*Final printed label on formulated Type B Medicated feed must bear a single concentration for each drug.  
The ratio between chlortetracycline and decoquinatate must be 44:1.

<sup>a</sup>Deccox® is the proprietary name of decoquinatate Type A medicated article (NADA 039-417).

<sup>b</sup>Pennchlor® is the proprietary name of chlortetracycline Type A medicated article (NADA 138-935).