

**Oxytetracycline Sheep Supplement
Type B Medicated Feed**

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For use in manufacturing medicated sheep feed.

ACTIVE DRUG INGREDIENT

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to *2-20 grams oxytetracycline hydrochloride/lb.

GUARANTEED ANALYSIS

Crude Protein, (Min).....	XX.XX %
NPN, (Max) ¹	X.XX%
Crude Fat, (Min).....	X.XX%
Crude Fiber, (Max).....	X.XX%
Calcium, (Min) ¹	X.XX%
Calcium, (Max) ¹	X.XX%
Phosphorus, (Min) ¹	X.XX%
Salt, (Min) ¹	X.XX%
Salt, (Max) ¹	X.XX%
Sodium, (Min) ²	X.XX%
Sodium, (Max) ²	X.XX%
Copper, (Min) ³	ppm
Copper, (Max) ³	ppm
Selenium, (Min) ¹	ppm
Vitamin A, (Min) ⁴	____IU/Lb

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³If added, or if total copper exceeds 20 ppm

⁴Other than precursor of Vitamin A, if added

INGREDIENTS

Ingredients as defined by AAFCO

MIXING DIRECTIONS

Mix this Type B medicated feed with non-medicated feed ingredients to prepare complete Type C medicated feed as described below. To ensure adequate mixing, an intermediate blending step should be used prior to manufacturing a complete feed.

Indications for Use

Oxytetracycline in
Type C medicated feed

SHEEP

Mix an amount of Type B to provide approximately

10 mg/lb body weight/day

10 mg/lb body weight/day varying with body weight and feed consumption with non-medicated feed ingredients to make the daily complete feed for treatment of bacterial enteritis caused by Escherichia coli and bacterial pneumonia caused by Pasteurella multocida susceptible to oxytetracycline to be fed continuously for 7-14 days.

RESIDUE WARNING: 5-day withdrawal period at 10 mg/lb dosage.

*The medicated feed label must state a single drug concentration.

Net Weight _____ lb (_____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000