Oxytetracycline Calf Ration for Treatment of Bacterial Enteritis Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For treatment of bacterial enteritis caused by <u>Escherichia coli</u> susceptible to oxytetracycline in calves for use in calf milk replacers or starter feeds.

ACTIVE DRUG INGREDIENT Oxytetracycline hydrochloride. 2000 g/ton* **GUARANTEED ANALYSIS** Crude Protein, (Min)..... XX.XX % NPN, (Max) ¹..... X.XX% X.XX% Crude Fiber, (Max)..... X.XX% Calcium, (Min) ¹..... X.XX% Calcium, (Max) ¹..... X.XX% Phosphorus, (Min) ¹..... X.XX% Salt, (Min) ¹..... X.XX% Salt, (Max) ¹..... X.XX%

Sodium, (Min) ².....

Sodium, (Max) ².....

Potassium, $(Min)^1$

Vitamin A, (Min)³.....

INGREDIENTS

X.XX%

X.XX%

X.XX%

____IU/Lb

Ingredients as defined by AAFCO

DIRECTIONS FOR USE

Feed continuously for 7-14 days as the sole ration in milk replacers or starter feeds.

RESIDUE WARNING: 5-day withdrawal period at 10 mg/lb dosage.

*This drug concentration will provide 10 mg/lb body weight/day when fed at a rate of 1.0 lb/100 lbs body weight/day.

Net Weight _____ lb (____ kg)

Manufactured by
Blue Bird Feed Company
Blue Bird, MD 00000

¹If added

²Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

³Other than precursors of Vitamin A, if added