

**Oxytetracycline Turkey Ration for Control of Complicating Bacterial Organisms Associated with  
Bluecomb  
Type C Medicated Feed**

**CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

For control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to oxytetracycline.

**ACTIVE DRUG INGREDIENT**

Oxytetracycline (from oxytetracycline dihydrate base) equivalent to \_\_\_\_\_ grams\* oxytetracycline hydrochloride/ton to provide approximately 25 mg/lb body weight/day varying with body weight and feed consumption.

**GUARANTEED ANALYSIS**

Crude Protein, (Min). . . . .	XX.XX %
Lysine, (Min). . . . .	X.XX%
Methionine, (Min). . . . .	X.XX%
Crude Fat, (Min). . . . .	X.XX%
Crude Fiber, (Max). . . . .	X.XX%
Calcium, (Min). . . . .	X.XX%
Calcium, (Max). . . . .	X.XX%
Phosphorus, (Min). . . . .	X.XX%
Salt, (Min) <sup>1</sup> . . . . .	X.XX%
Salt, (Max) <sup>1</sup> . . . . .	X.XX%
Sodium, (Min) <sup>2</sup> . . . . .	X.XX%
Sodium, (Max) <sup>2</sup> . . . . .	X.XX%

<sup>1</sup>If added

<sup>2</sup>Should be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee

**INGREDIENTS**

Ingredients as defined by AAFCO

**DIRECTIONS FOR USE**

Feed continuously for 7-14 days as the sole ration.

**RESIDUE WARNING: Do not feed to turkeys producing eggs for human consumption.  
Zero-day withdrawal period.**

\*The medicated feed label must state a single drug concentration.

Net Weight \_\_\_\_\_ lb (\_\_\_\_\_ kg)

Manufactured by  
Blue Bird Feed Company  
Blue Bird, MD 00000